

CONTENTS

1	INTRODUCTION	1
2	PREVIOUS INVESTIGATIONS	3
3	METHODS	4
4	REEF MORPHOLOGY AND LITHOFACIES	8
5	REEF PETROLOGY	15
6	REEF LITHOLOGIC ASSEMBLAGES	22
7	CEMENTATION	24
8	CORAL GROWTH AND REEF ACCRETION RATES	26
9	DISCUSSION	30
10	CONCLUSIONS	41
11	REFERENCES	42

LIST OF FIGURES

Figure 1.	General location of the Cape Tribulation area	2
Figure 2.	Location of Emmagen, Rykers and South Myall Reefs	2
Figure 3.	Surface lithofacies, Cape Tribulation reefs	6
Figure 4.	Surveyed reef surfaces, Emmagen to Rykers	9
Figure 5.	Surveyed reef surfaces, South Tribulation to Noah's Reef	10
Figure 6.	Portable drilling rig on Rykers Reef	11
Figure 7.	Terrigenous cobbles, Rykers Reef. Mangrove is about 1.5 m tall	11
Figure 8.	Back reef pool, Rykers Reef. Low elevation areas of the back reef are often characterised by the presence of <i>Sargassum</i> sp	13
Figure 9.	High energy conditions on the outer reef crest restrict growth to durable turf algae, South Cape Tribulation Reef	13
Figure 10.	Reef crest of the narrow fringing reef crest south of Emmagen Creek, pictured during extremely low Spring tides	14
Figure 11.	Subtidal branching and foliose living coral, approximately 2 m below water surface at Rykers Reef. Sediment traps used in a parallel study can be seen at the reef base.	14
Figure 12.	Borehole logs R1, R2, R3 and R4, Rykers Reef	18
Figure 13.	Borehole logs M1, M2 and M3, South Myall Reef	20
Figure 14.	Borehole logs, E1 and E2, Emmagen Reef	21
Figure 15.	Cross sections of Emmagen, Rykers and South Myall Reefs	23
Figure 16.	Acicular aragonite cement lining the pores of <i>Porites</i> sp sample M1-5B from a depth of 8.3 m below reef surface (7.2 m below Cairns Port Datum)	25
Figure 17.	Positive image radiograph showing growth bands of coral core samples M3-2A, M3-2B and M3-3C, South Myall Reef. Samples were obtained from a depth of 1.3-2.0 m below the surface (0.4-1.6 m below Cairns Port Datum). Core width is 4.5 cm.	27
Figure 18.	Vertical accretion rates (mm/yr) plotted against depth of water and time of growth (m) for fringing, mid-shelf and outer shelf reefs of the Great Barrier Reef (from dated cores of PJ Davies <i>et al</i> , BMR and D Hopley <i>et al</i> , JCU) ...	31
Figure 19.	Radiocarbon ages from reef cores. Ages are in C ¹⁴ years before present	35
Figure 20.	Reef accretion in relation to sea level rise, Emmagen, Rykers and South Myall Reefs	36
Figure 21.	Classification of fringing reef development based on Great Barrier Reef examples.	39

LIST OF TABLES

Table 1.	Coral species identified from borehole samples	16
Table 2.	Cape Tribulation fringing reefs - coral growth rates determined by measuring radiographs of reef cores	28
Table 3.	Reef vertical accretion rates, Cape Tribulation reefs, determined by C14 age-dating . .	28
Table 4.	Cape Tribulation fringing reefs - coral growth rates determined by measuring radiographs of reef cores (samples are listed in order of increasing depth)	29
Table 5.	Coring investigations of fringing reefs in the Great Barrier Reef system	30
Table 6.	Cape Tribulation mean coral growth rates determined by measuring radiographs of reef cores	33
Table 7.	Holocene vegetation and climate change on the Atherton Tableland (from Kershaw) related to stages in reef development, Cape Tribulation	38