

## APPENDIX

**Table 2.** Number of intertidal and sub-tidal sites used to calculate mean above-ground biomass at each survey transect, north and south, of Oyster Point in November 1995, December 1997 and November 1998.

Transect distance (km)	Number of sites															
	North								South							
	Intertidal				Sub-tidal				Intertidal				Sub-tidal			
	1995	1997	1998	1999	1995	1997	1998	1999	1995	1997	1998	1999	1995	1997	1998	1999
0.0	0	1	1	0	1	3	2	2								
0.1	1	3	3	2	4	2	4	1	0	0	0	0	1	0	0	0
0.2	6	4	5	3	7	3	4	3	3	0	1	0	3	0	0	0
0.3	3	5	3	5	6	4	4	5	1	0	2	5	2	1	3	1
0.4	3	6	6	2	7	8	4	8	7	10	8	6	2	1	3	3
0.5	3	4	4	3	10	8	3	7	8	10	6	3	3	3	4	2
0.6	2	4	4	3	10	7	5	8	9	7	14	4	3	1	1	1
0.7	2	2	9	2	10	6	6	8	6	5	6	4	7	3	3	6
0.8	2	4	4	3	19	7	7	10	3	3	3	4	6	4	2	7
0.9	4	4	5	3	18	7	4	9	2	2	5	2	8	8	2	10
1.0	4	3	4	3	15	11	8	11	3	4	5	3	7	5	7	5
1.5	3	5	5	3	18	11	9	13	6	3	2	2	13	7	7	7
2.0	3	6	3	2	14	17	11	14	3	4	3	3	5	9	6	8

**Table 3.** Results of regressions of each diver's seagrass biomass estimation with harvested above-ground biomass (g DW m<sup>-2</sup>).

Diver	November 1995			December 1997			November 1998			December 1999		
	r <sup>2</sup>	F	P	r <sup>2</sup>	F	p	r <sup>2</sup>	F	p	r <sup>2</sup>	F	p
G. Chisholm	0.97	117.54	0.001									
R.G. Coles	0.92	48.56	0.002	0.79	37.81	<0.001						
L. Makey	0.88	29.32	0.006									
P.A. Daniel				0.77	33.14	<0.001	0.86	30.80	0.003			
W.L. Lee Long	0.99	411.43	<0.001	0.98	207.25	<0.001	0.88	35.84	0.002	0.98	157.71	<0.001
L.J. McKenzie	0.97	191.02	<0.001									
C.A. Roder				0.96	249.70	<0.001						
A.J. Roelofs	0.97	156.84	<0.001	0.99	244.43	<0.001	0.90	47.26	0.001	0.86	25.00	0.008
R. Thomas										0.82	18.72	0.012
T. Curruthers*										0.95	80.15	<0.001

\* Analysis performed on Ln (x+1) data.