

TABLE OF CONTENTS

	Page
1. INTRODUCTION	1
2. CAIRNS AREA	2
3. INNISFAIL AREA	3
4. TOWNSVILLE AREA	5
5. MACKAY AREA	7
6. CAPRICORN-BUNKER AND LADY ELLIOTT ISLAND AREA	9
7. CHANGES IN CATCH WITH LATITUDE	12
8. DISCUSSION	14
9. ACKNOWLEDGMENTS	16

LIST OF APPENDICES

I DISTRIBUTION LIST	17
II DOCUMENT CONTROL DATA PAGE	18

LIST OF FIGURES

1. CATCH vs DISTANCE FROM CAIRNS	2
2. CATCH AND MEAN FISH WEIGHT OVER TIME (INNISFAIL)	3
3. CATCH vs DISTANCE FROM INNISFAIL	4
4. CATCH AND MEAN FISH WEIGHT OVER TIME (TOWNSVILLE)	5
5. CORAL TROUT AND RED SWEETLIP AS PERCENTAGE OF CATCH (TOWNSVILLE)	6
6. CATCH OF MAJOR SPECIES OVER TIME (TOWNSVILLE)	7
7. CATCH AND MEAN FISH WEIGHT OVER TIME (MACKAY)	8
8. CORAL TROUT AS PERCENTAGE OF CATCH (MACKAY)	9
9. CATCH AND MEAN FISH WEIGHT OVER TIME (CAPRICORN-BUNKER)	10

	Page
10. MEAN WEIGHT OF LARGEST OF SPECIES CAUGHT EACH TRIP FOR CORAL TROUT, RED EMPEROR AND RED SWEETLIP (CAPRICORN-BUNKER)	11
11. CATCH AND MEAN FISH WEIGHT vs DISTANCE FROM NEAREST PORT (CAPRICORN-BUNKER AND CONTINENTAL SHELF)	12
12. CATCH AND MEAN FISH WEIGHT BY LATITUDE	13