

REFERENCES

- Bulthuis, D.A. (1987). Effects of temperature on photosynthesis and growth of seagrasses. *Aquatic Botany* **27**, 27-41.
- Coles, R.G. and Lee Long, W.J. (1985). Juvenile prawn biology and the distribution of seagrass prawn nursery grounds in the southeastern Gulf of Carpentaria. In 'Second Australian National Prawn Seminar, Cleveland.' (Eds P. P. Rothlisberg, D. J. Hill and D. J. Staples.) pp. 55-60. (NPS2: Cleveland, Australia.).
- Coles, R.G., Lee Long, W.J., Squire, B.A., Squire, L.C. and Bibby, J.M. (1987a). Distribution of seagrasses and associated juvenile commercial penaeid prawns in north-eastern Queensland waters. *Australian Journal Marine and Freshwater Research* **38**, 103-119.
- Coles, R.G. Mellors, J.E., Bibby, J. and Squire, B. (1987b). Seagrass beds and juvenile prawn nursery grounds between Bowen and Water Park Point, A report to the Great Barrier Reef Marine Park Authority. Queensland Department of Primary Industries Information Series QI 87021. (Queensland Gov, Brisbane.) 54 pp.
- Coles, R.G., Lee Long, W.J., Watson, R.A. and Derbyshire, K.J. (1993). Distribution of seagrasses, and their fish and penaeid prawn communities, in Cairns Harbour, a tropical estuary, northern Queensland, Australia. In 'Tropical Seagrass Ecosystems; Structure and Dynamics in the Indo-West Pacific.' *Australian Journal of Marine and Freshwater Research* **44**, 193-210.
- Commonwealth Commission of Inquiry (1994) "Shoalwater Bay, Capricornia Coast, Queensland - Interim Report No. 2, March 1994." (Aust. Gov. Publ. Serv.: Canberra). 503 pp.
- Dall, W. (1957). Observations on the biology of the Greentail prawn *Metapenaeus masterii* (Haswell) (Crustacea, Decapoda: Penaeidae). *Australian Journal of Marine and Freshwater Research* **9**, 111-134.
- Dennison, W.C. (1987). Effects of light on seagrass photosynthesis, growth and depth distribution. *Aquatic Botany* **27**, 15-27.
- Dennison, W. C. and Kirkman, H. (1996). Seagrass survival model. In: Seagrass Biology: Proceedings of an International Workshop, Rottnest Island, Western Australia, 25-29 Jan 1996. (Eds: Kuo, J.; Phillips, R. C; Walker, D. I; Kirkman, H.) Faculty of Sciences, University of Western Australia, Nedlands, Perth, Australia, 341-344.
- Dennison, W.C., Orth, R.J., Moore, K.A., Stevenson, J.C., Carter, V., Kollar, S., Bergstrom, P.W., and Batiuk, R.A. (1993). Assessing water quality with submersed aquatic vegetation: Habitat requirements as barometers of Chesapeake Bay health. *BioScience* **43**(2), 86-94.
- Downing, J.A. and Anderson, M.R. (1985). Estimating the standing biomass of aquatic macrophytes. *Canadian Journal of Fisheries and Aquatic Sciences*. **42**, 1860-1869.
- Erftemeijer, P.L.A. (1994). Differences in nutrient concentrations and resources between seagrass communities on carbonate and terrigenous sediments in South Sulawesi, Indonesia. *Bulletin of Marine Science* **54**(2), 403-419.
- Fitzsimmons, L (Ed.) (1996): Status of fisheries and fisheries resources in Shoalwater Bay, A report to the Great Barrier Reef Marine Park Authority. Queensland Department of Primary Industries, Southern Fisheries Centre, Beach Road, Deception Bay, Qld. 100 pp.
- Great Barrier Reef Marine Park Authority. (1997). 'Shoalwater Bay (Dugong) plan of management'. No. SB01/97 (Great Barrier Reef Marine Park Authority, Townsville) 8 pp.
- Grey, D.L., Dall, W., and Baker, A. (1983). 'A guide to the Australian Penaeid prawns'. (Northern Territory Government Printing, Darwin) 140 pp.

- Hyland, S.J., Courtney, A.J., and Butler, C.T. (1989). 'Distribution of seagrass in the Moreton Region from Coolangatta to Noosa'. Queensland Department of Primary Industries Information Series #QI89010 (Queensland Department of Primary Industries, Brisbane) 42 pp.
- Jackson, J.B.C., Cubit, J.D., Keller, B.D., Batista, V., Burns, K., Caffey, H.M., Caldwell, R.L., Garrity, S.D., Getter, C.D., Gonzalez, C., Guzman, H.M., Kaufmann, K.W., Knap, A.H., Levings, S.C., Marshall, M.J., Steger, R., Thompson, R.C., and Weil, E. (1989). Ecological effects of a major oil spill on Panamanian coastal marine communities. *Science* **243**, 37-44.
- Kirkman, H. and Kuo, J. (1990). Pattern and process in southern Western Australian seagrasses. *Aquatic Botany*. **37**, 367-382.
- Kuo, J. and McComb, A.J. (1989). Seagrass taxonomy, structure and development. In 'Biology of seagrasses: A treatise on the biology of seagrasses with special reference to the Australian region.' (Eds A.W.D. Larkum, A.J. McComb and S.A. Shepherd) Chpt.2. pp.6-73. (Elsevier, New York).
- Larkum, A.W.D., McComb, A.J., and Shepherd, S.A. (1989). 'Biology of Seagrasses. A Treatise on the biology of seagrasses with special reference to the Australian region'. (Eds.) (Elsevier Science Publ., Amsterdam, New York.) 841 pp.
- Lanyon, J., Limpus, C.J. and Marsh, H. (1989). Dugongs and turtles: grazers in the seagrass system. In 'Biology of seagrasses: A treatise on the biology of seagrasses with special reference to the Australian region.' (Eds A.W.D. Larkum, A.J. McComb and S.A. Shepherd) Chpt.18. pp.610-634. (Elsevier, New York).
- Lee Long, W.J., Mellors, J.E. and Coles, R.G. (1993). Seagrasses between Cape York and Hervey Bay, Queensland, Australia. *Australian Journal of Marine and Freshwater Research* **44**(1), 19-31.
- Lee Long, W.J., Mellors, J.E. and Coles, R.G. (1994). Corrigendum to Seagrasses between Cape York and Hervey Bay, Queensland, Australia. *Australian Journal of Marine and Freshwater Research* **45**(4), 727.
- Lee Long, W.J., Coles, R.G. and McKenzie, L.J. (1996). Deepwater seagrass in northeastern Australia - how deep, how meaningful?. In 'Seagrass Biology: Proceedings of an International Workshop, Rottnest Island, Western Australia, 25-29 January 1996.' (Eds J. Kuo, R.C. Phillips, D.I. Walker and H. Kirkman) pp. 345-350. (Western Australian Museum, Perth).
- Long, B.G., Skewes, I.D., and Poiner, I.R. (1994). An efficient method for estimating seagrass biomass. *Aquatic Botany*. **47**(3-4), 277-292.
- Marsh, H., Corkeron, P., Lawler, I.R., Lanyon, J., and Preen, A.R. (1995). 'The status of the dugong in the southern Great Barrier Reef Marine Park'. Unpublished report the Great Barrier Reef Marine Park Authority (TESAG, Townsville) 84 pp.
- McKenzie, L.J. (1994). Seasonal changes in biomass and shoot characteristics of a *Zostera capricorni* Aschers. dominant meadow in Cairns Harbour, northern Queensland. *Australian Journal of Marine and Freshwater Research* **45**, 1337-1352.
- Mellors, J.E. (1991). An evaluation of a rapid visual technique for estimating seagrass biomass. *Aquatic Botany* **42**, 67-73.
- Mellors, J.E., Marsh, H., and Coles, R.G. (1993). Intra-annual changes in seagrass standing crop, Green Island, northern Queensland. *Australian Journal of Marine and Freshwater Research* **44**, 33-42.
- Orth, R.J. (1977). Effect of nutrient enrichment on growth of eelgrass *Zostera marina* in the Chesapeake Bay, Virginia, USA. *Marine Biology*. **44**, 187-194.

- Orth, R.J. and van Montfrans, J. (1987). Utilisation of a seagrass meadow and tidal marsh creek by blue crabs *Callinectes sapidus*. I. Seasonal and annual variations in abundance with emphasis on post-settlement juveniles. *Marine Ecology Progress Series* **41**, 283-294.
- Preen, A.R. and Marsh, H. (1995). Response of Dugongs to Large-scale Loss of Seagrass from Hervey Bay, Queensland, Australia. *Wildlife Research*. **22**, 507-19.
- Poiner, I.R., Walker, D.I., and Coles, R.G. (1989). Regional studies - seagrasses of tropical Australia. In. 'Biology of Seagrasses: A treatise on the biology of seagrasses with special reference to the Australian region'. (Eds. A. W. D. Larkum, A. J. McComb, and S. A. Shepherd.) chpt. 10, pp. 279-296. (Elsevier: New York).
- Preen, A.R., Lee Long, W.J., and Coles, R.G. (1995). Flood and cyclone related loss, and partial recovery, of more than 1,000 km² of seagrass in Hervey Bay, Queensland, Australia. *Aquatic Botany* **52**, 3-17.
- Pringle, A.W. (1989). 'The history of dredging in Cleveland Bay, Queensland and its effect on sediment movement and on the growth of mangroves, corals and seagrass'. Research Publication (GBRMPA, Townsville) 177 pp.
- Queensland Department of Transport (1995) "Queensland's Official Tide Tables & Boating Guide". (QDT: Brisbane) 216 pp.
- Shepherd, S.A., McComb, A.J., Bulthuis, D.A., Neverauskas, V., Steffensen, D.A. and West, R. (1989). Decline of seagrasses. In. 'Biology of Seagrasses: A treatise on the biology of seagrasses with special reference to the Australian region'. (Eds. A. W. D. Larkum, A. J. McComb, and S. A. Shepherd.) chpt. 12, pp. 346-387. (Elsevier: New York).
- Short, F.T. (1987). Effects of sediment nutrients on seagrasses: Literature review and mesocosm experiment. *Aquatic Botany* **27**, 41-57.
- Sokal, R.R. and Rolf, F.J. (1987). 'Introduction to Biostatistics'. (W.H. Freeman, New York) 363pp.
- Thresher, R.E. and Gunn, J.S. (1986). Comparative analysis of visual census techniques for highly mobile, reef associated piscivores (Carangidae). *Environmental Biology of Fishes*. **17**(2), 93-116.
- Walker, D.I. (1989). 'Methods for monitoring seagrass habitat'. Victorian Institute of Marine Science Working Paper 18 (Victorian Institute of Marine Science, Queenscliff) 21 pp.
- Watson, R.A., Coles, R.G. and Lee Long, W.J. (1993). Simulation estimates of annual yield, and landed value for commercial penaeid prawns from a tropical estuary, northern Queensland, Australia. *Australian Journal of Marine and Freshwater Research* **44**(1), 211-219.
- Williams, S.L. and Dennison, W.C. (1990). Light availability and diurnal growth of a green macro-alga (*Caulerpa cupressoides*) and a seagrass (*Halophila decipiens*). *Marine Biology* **106**(3), 437-443.